

# FRACTAL BOUNDARY VALUE PROBLEMS FOR INTEGRAL AND DIFFERENTIAL EQUATIONS WITH LOCAL FRACTIONAL OPERATORS

Xiao-Jun YANG<sup>a</sup>, Dumitru BALEANU<sup>b,c,d</sup>, Mihailo P. LAZAREVIĆ<sup>e</sup>,  
Milan S. CAJIĆ<sup>f,\*</sup>

<sup>a</sup>Department of Mathematics and Mechanics, China University of Mining and Technology, Xuzhou, 221008, China

<sup>b</sup>Department of Mathematics and Computer Sciences, Faculty of Arts and Sciences, Cankaya University, 06530, Ankara, Turkey

<sup>c</sup>Department of Chemical and Materials Engineering, Faculty of Engineering, King Abdulaziz University, P.O. Box 80204, Jeddah, 21589, Saudi Arabia

<sup>d</sup>Institute of Space Sciences, Magurele-Bucharest, Romania

<sup>e</sup>Faculty of Mechanical Engineering, Kraljice Marije 16, 11120 Belgrade, Serbia

<sup>f</sup>Mathematical Institute SANU, Kneza Mihaila 36, 11001 Belgrade, Serbia

*In the present paper we investigate the fractal boundary value problems for the Fredholm\Volterra integral equations, heat conduction and wave equations by using the local fractional decomposition method. The operator is described by the local fractional operators. The four illustrative examples are given to elaborate the accuracy and reliability of the obtained results.*

*Keywords: local fractional decomposition method, heat conduction equations, integral equations, wave equations, boundary value problems*

## 1. Introduction

Fractals are used in many engineering applications such as porous media modelling, nano fluids, fracture mechanics and many other applications in nanoscale [1,2] where various transport phenomena cannot be described by smooth continuum approach and need the fractal nature of the objects to be taken into account. For the transport phenomena performed in fractal objects the local temperature depends on the fractal dimensions where adequate physical results can be achieved by the application of local fractional models and relevant solution approaches. For example, in [3] heat diffusion problem is analyzed in fractal geometry cooling surface and in [4] fractal heat conduction problem is solved by using the local fractional variation iteration method.

The decomposition method introduced by Adomian [5, 6] was broadly applied to solve a wide class of linear and nonlinear problems described by ordinary, partial differential equations and integral equations[7-9]. There are exist some other analytical methods widely applied to solve non-linear problems, among them: the variational iteration method [10, 11], the homotopy perturbation method [12, 13], the heat-balance integral method [14, 15], the complex transform method [16, 17], the

---

\* Corresponding author. Tel.: +381 62 83 68 505 (Mobile); Fax: +381 11 21 86 105

E-mail adresses: mcajic@mi.sanu.ac.rs (M. Cajić), caji84@gmail.com

homotopy analysis method [18], the fractional sub-equation method [19] and the fractional variational iteration method [20-22] and more details seen in [23].

The local fractional derivative is a relatively new tool to investigate local fractal behaviors of differential equations with fractal conditions [24-28]. Several works have been published on efficient methods for solving such local fractional differential equations with non-differential functions [29, 31, 39, 40]. The integral equations via local fractional calculus theory [27, 28] were first proposed in [29] and developed in the case of local fractional Fredholm and Volterra integral equations [30]. For the differential and integral equations with local fractional derivative and integral operators, there are exist some analytical methods, such as: the local fractional variational iteration method [31, 32], the local fractional decomposition method [33, 34], the local fractional Fourier series method [4, 35], the local fractional Fourier transform method [12, 36], the local fractional Laplace transform method [12], the local fractional Z transform method [37], etc.

Taking into account the existing background in solution of fractal boundary problems the present communication addresses analytical solutions performed by the local fractional decomposition method to two problems: the local fractional integral equations and the local fractional wave equation. The manuscript is organized as follows: Section 2 is devoted to the local fractional decomposition method. The four illustrative examples are shown in Section 3. Finally, conclusions are given in Section 4.

## 2. Analytical method

For seek of clarity of the explanation, the local fractional decomposition method will be briefly outlined. For integral equations a compact recurrence scheme has been developed [7-30]. The initial (zeroth) approximation in this case is

$$u_0(x) = f(x), \quad (1)$$

and

$$u_{n+1}(x) = \frac{\lambda^\alpha}{\Gamma(1+\alpha)} \int_a^b K(x,t) u_n(x) (dt)^\alpha, n \in N. \quad (2)$$

Local fractional integral of  $f(x)$  of order  $\alpha$  ( $0 < \alpha \leq 1$ ) in the interval  $[a, b]$  is denoted by [23-35]

$${}_a I_b^{(\alpha)} f(x) = \frac{1}{\Gamma(1+\alpha)} \int_a^b f(t) (dt)^\alpha = \frac{1}{\Gamma(1+\alpha)} \lim_{\Delta t \rightarrow 0} \sum_{j=0}^{j=N-1} f(t_j) (\Delta t_j)^\alpha, \quad (3)$$

with a partition of the interval  $[a, b]$ , which is defined through  $(t_j, t_{j+1})$  with  $\Delta t_j = t_{j+1} - t_j$ ,  $\Delta t = \max\{\Delta t_1, \Delta t_2, \Delta t_j, \dots\}$ ,  $j = 0, \dots, N-1$ ,  $t_0 = a$  and  $t_N = b$ .

Hence, we can determine a few terms in the series such as  $u(x) = \sum_{n=0}^{\infty} u_n(x)$  by truncating the series at certain term. The local fractional Volterra integral equation is written in the form [26, 30]

$$u(x) = f(x) + \frac{\lambda^\alpha}{\Gamma(1+\alpha)} \int_0^x K(x,t) u(t) (dt)^\alpha, \quad (4)$$

and hence

$$u_0(x) = f(x). \quad (5)$$

Substituting  $u(x) = \sum_{n=0}^{\infty} u_n(x)$  into eq. (4) implies

$$\sum_{n=0}^{\infty} u_n(x) = f(x) + \frac{\lambda^\alpha}{\Gamma(1+\alpha)} \int_0^x K(x,t) \left\{ \sum_{n=0}^{\infty} u_n(x) \right\} (dt)^\alpha. \quad (6)$$

The components  $u_0(x), u_1(x), u_2(x), \dots, u_n(x), \dots$  of the function  $u(x)$  can be completely determined if we set

$$\begin{aligned} u_0(x) &= f(x) \\ u_1(x) &= \frac{\lambda^\alpha}{\Gamma(1+\alpha)} \int_0^x K(x,t) u_0(x) (dt)^\alpha \\ u_2(x) &= \frac{\lambda^\alpha}{\Gamma(1+\alpha)} \int_0^x K(x,t) u_1(x) (dt)^\alpha \\ &\dots \\ u_n(x) &= \frac{\lambda^\alpha}{\Gamma(1+\alpha)} \int_0^x K(x,t) u_{n-1}(x) (dt)^\alpha \end{aligned} \quad (7)$$

and so on. The set of above equations can be written in compact recurrence scheme as

$$u_0(x) = f(x), \quad (8)$$

and

$$u_{n+1}(x) = \frac{\lambda^\alpha}{\Gamma(1+\alpha)} \int_0^x K(x,t) u_n(x) (dt)^\alpha, n \in N. \quad (9)$$

Hence, we give the local fractional series solution  $u(x) = \sum_{n=0}^{\infty} u_n(x)$ . The above processes were discussed in [37, 38].

Let us consider the local fractional Volterra integral equation given through

$$u(x) = f(x) + \frac{\lambda^\alpha}{\Gamma(1+\alpha)} \int_0^a K(x,t)u(t)(dt)^\alpha, \quad (10)$$

where  $a$  is a constant. We can set up on an iteration formula based on the decomposition method

$$u_{n+1}(x) = \frac{\lambda^\alpha}{\Gamma(1+\alpha)} \int_a^b K(x,t)u_n(x)(dt)^\alpha, n \in N. \quad (11)$$

An initial condition can be expressed as

$$u_0(x) = f(x). \quad (12)$$

Thus, we arrive at a local fractional series solution, namely:

$$u(x) = \sum_{n=0}^{\infty} u_n(x). \quad (13)$$

Moreover, we can write the general local fractional differential equation in a local fractional differential operator form

$$L_x^{(2\alpha)}u(x) + R_x^\alpha u(x) = f(x), \quad (14)$$

In eq. (14)  $L_x^{(2\alpha)}$  is local fractional  $2\alpha^{\text{th}}$  order differential operator [27, 28], which by the definition reads

$$L_x^{(2\alpha)}s(x) = \frac{d^\alpha}{dx^\alpha} \left[ \frac{d^\alpha}{dx^\alpha} s(x) \right], \quad (15)$$

and

$$R_x^\alpha s(x) = \frac{d^\alpha s(x)}{dx^\alpha} \Big|_{x=x_0} = \lim_{x \rightarrow x_0} \frac{\Delta^\alpha (s(x) - s(x_0))}{(x - x_0)^\alpha}, \quad (16)$$

is local fractional  $\alpha^{\text{th}}$  order differential operator ( $0 < \alpha \leq 1$ ) and  $s(x)$  is local fractional continuous

[27-33].

Applying the inverse operator  $L_x^{(-2\alpha)}$  to both sides of (14) yields

$$L_x^{(-2\alpha)} L_x^{(2\alpha)} u(x) = -L_x^{(-2\alpha)} R_x^\alpha u(x) + L_x^{(-2\alpha)} f(x). \quad (17)$$

If the inverse differential operator  $L_x^{(-2\alpha)}$  exists, according to the local fractional decomposition method mentioned above, we have

$$\begin{cases} u_{n+1}(x) = -L_x^{(-2\alpha)} R_x^\alpha u_n(x), \\ u_0(x) = r(x) \end{cases}, \quad (18)$$

where  $r(x) = L_x^{(-2\alpha)} f(x)$  and

$$L_x^{(-2\alpha)} u(x) = \left( \frac{1}{\Gamma(1+\alpha)} \right)^2 \int_0^x \int_0^{t_2} u(t_1) (dt_1)^\alpha (dt_2)^\alpha. \quad (19)$$

Proof of the existence of the inverse differential operator  $L_x^{(-2\alpha)}$  as local fractional integral operator one can find in [28]. Finally, we can find a solution in the form

$$u(x) = \sum_{n=0}^{\infty} u_n(x). \quad (20)$$

Hence, we can obtain that [27-28, 34] the following condition is obeyed

$$|f(x) - f(x_0)| < \varepsilon^\alpha, \quad (21)$$

where fractal dimension of  $f(x)$  is equal to  $\alpha$  for any  $x \in (a, b)$ .

### 3. Illustrative examples

Several illustrative examples demonstrating the efficiency of the of the suggested local fractional decomposition method are present next

#### Example 1.

First, we solve the local fractional Fredholm integral equation

$$u(x) = \frac{x^\alpha}{\Gamma(1+\alpha)} + \frac{1}{\Gamma(1+\alpha)} \int_0^1 \frac{(t-1)^\alpha}{\Gamma(1+\alpha)} u(t) (dt)^\alpha. \quad (22)$$

The zeroth approximation is as follows:

$$u_0(x) = \frac{x^\alpha}{\Gamma(1+\alpha)}. \quad (23)$$

The first approximation is expressed through

$$\begin{aligned} u_1(x) &= \frac{1}{\Gamma(1+\alpha)} \int_0^1 \frac{(t-1)^\alpha}{\Gamma(1+\alpha)} \frac{t^\alpha}{\Gamma(1+\alpha)} (dt)^\alpha. \\ &= -\frac{1}{\Gamma(1+3\alpha)} \end{aligned} \quad (24)$$

Proceeding in this manner, we can find the second approximation as

$$\begin{aligned} u_2(x) &= \frac{1}{\Gamma(1+\alpha)} \int_0^1 \frac{(t-1)^\alpha}{\Gamma(1+\alpha)} u_1(t) (dt)^\alpha \\ &= -\frac{1}{\Gamma(1+\alpha)} \int_0^1 \frac{(t-1)^\alpha}{\Gamma(1+\alpha)} \frac{1}{\Gamma(1+3\alpha)} (dt)^\alpha. \\ &= \frac{1}{\Gamma(1+3\alpha)} \frac{1}{\Gamma(1+2\alpha)} \end{aligned} \quad (25)$$

Finally, we get

$$u_n = -\frac{1}{\Gamma(1+3\alpha)} \left( \frac{1}{\Gamma(1+2\alpha)} \right)^{n-1}, n \in N. \quad (26)$$

Hence, we arrive at the solution

$$\begin{aligned}
& u(x) \\
&= \lim_{n \rightarrow \infty} \sum_{n=0}^{\infty} u_n(x) \\
&= \frac{x^\alpha}{\Gamma(1+\alpha)} + \lim_{n \rightarrow \infty} \left( -\frac{1}{\Gamma(1+3\alpha)} \sum_{n=1}^{\infty} \left( \frac{1}{\Gamma(1+2\alpha)} \right)^{n-1} \right) \\
&= \frac{x^\alpha}{\Gamma(1+\alpha)} - \frac{1}{\Gamma(1+3\alpha)} \frac{\frac{1}{\Gamma(1+2\alpha)}}{1 - \frac{1}{\Gamma(1+2\alpha)}} \\
&= \frac{x^\alpha}{\Gamma(1+\alpha)} - \frac{1}{\Gamma(1+3\alpha)(\Gamma(1+2\alpha)-1)}
\end{aligned} \tag{27}$$

If  $n \rightarrow \infty$ .

**Example 2.**

Similarly, for the solution of the local fractional Volterra integral equation

$$u(x) = 1 + \frac{1}{\Gamma(1+\alpha)} \int_0^x \frac{(x-t)^\alpha}{\Gamma(1+\alpha)} u(t) (dt)^\alpha . \tag{28}$$

the zeroth approximation is suggested as

$$u_0(x) = 1. \tag{29}$$

Then, the second approximation is

$$\begin{aligned}
& u_1(x) \\
&= \frac{1}{\Gamma(1+\alpha)} \int_0^x \frac{(x-t)^\alpha}{\Gamma(1+\alpha)} (dt)^\alpha \\
&= \frac{x^{2\alpha}}{\Gamma(1+2\alpha)}
\end{aligned} \tag{30}$$

Proceeding in this manner, the third approximation is

$$\begin{aligned}
u_2(x) &= \frac{1}{\Gamma(1+\alpha)} \int_0^x \frac{(x-t)^\alpha}{\Gamma(1+\alpha)} u_1(t) (dt)^\alpha \\
&= \frac{1}{\Gamma(1+\alpha)} \int_0^x \frac{(x-t)^\alpha}{\Gamma(1+\alpha)} \left( \frac{t^{2\alpha}}{\Gamma(1+2\alpha)} \right) (dt)^\alpha \\
&= \frac{x^{4\alpha}}{\Gamma(1+4\alpha)}
\end{aligned} \tag{31}$$

Therefore, we have

$$u_n = \frac{x^{2n\alpha}}{\Gamma(1+2n\alpha)}, n \in N. \tag{32}$$

In conclusion, when  $n \rightarrow \infty$  we arrive at

$$\begin{aligned}
u(x) &= \lim_{n \rightarrow \infty} \sum_{n=0}^{\infty} u_n(x) \\
&= \lim_{n \rightarrow \infty} \left( \sum_{n=0}^{\infty} \frac{x^{2n\alpha}}{\Gamma(1+2n\alpha)} \right) \\
&= \cosh_\alpha(x^\alpha)
\end{aligned} \tag{33}$$

where  $\cosh_\alpha x^\alpha = \sum_{k=0}^{\infty} x^{2\alpha k} / \Gamma(1+2\alpha k)$  is a hyperbolic cosine function defined on a Cantor set [27-28, 34].

**Example 3.**

Let us consider the local fractional heat conduction equation with no heat generation in fractal media and dimensionless variables [4], which reads

$$\frac{\partial^\alpha u(x,t)}{\partial t^\alpha} = \frac{\partial^{2\alpha} u(x,t)}{\partial x^{2\alpha}}, \tag{34}$$

subject to the following fractal initial boundary conditions

$$u_x^{(\alpha)}(x,0) = 0, \quad u(x,0) = E_\alpha(x^\alpha), \quad (0 \leq x \leq l), \tag{35}$$

where in eq. (34)  $u(x,t) = T(x,t)$  is the temperature field.



Hence, the recurrence formula takes the form

$$\begin{cases} u_{n+1}(x, t) = L_t^{(-\alpha)} L_{xx}^{(2\alpha)} u_n(x, t) \\ u_0(x, t) = E_\alpha(x^\alpha) \end{cases}. \quad (36)$$

We can develop a solution in a form of local fractional series, namely

$$\begin{aligned} u(x, t) &= \lim_{i \rightarrow \infty} \sum_{i=0}^{\infty} u_i(x, t) \\ &= E_\alpha(x^\alpha) \lim_{i \rightarrow \infty} \left( \sum_{i=0}^n \frac{t^{(2i+1)\alpha}}{\Gamma(1+(2i+1)\alpha)} \right) \\ &= E_\alpha(x^\alpha) \sinh_\alpha(t^\alpha) \end{aligned} \quad (37)$$

where  $\sinh_\alpha t^\alpha = \sum_{k=0}^{\infty} t^{\alpha(2k+1)} / \Gamma(1+(2k+1)\alpha)$  is a hyperbolic cosine function defined on a Cantor set [27, 28, 34], and  $E_\alpha(x^\alpha) = \sum_{k=0}^{\infty} x^{k\alpha} / \Gamma(1+k\alpha)$  is the Mittag-Leffler function defined on a Cantor set [27, 28, 34-35].

**Example 4.**

Consider the local fractional wave equation [41]

$$\frac{\partial^{2\alpha} u(x, t)}{\partial t^{2\alpha}} = a^{2\alpha} \frac{\partial^{2\alpha} u(x, t)}{\partial x^{2\alpha}} + \frac{\partial^\alpha u(x, t)}{\partial x^\alpha}, \quad (38)$$

with fractal initial boundary conditions

$$u_t^{(\alpha)}(x, 0) = 0, \quad u(x, 0) = \frac{x^{2\alpha}}{\Gamma(1+2\alpha)} \quad (0 \leq x \leq l). \quad (39)$$

Hence, the structure of the recurrence formula is:

$$\begin{cases} u_{n+1}(x,t) = a^{2\alpha} L_t^{(-2\alpha)} L_{xx}^{(2\alpha)} u_n(x,t) + L_t^{(-2\alpha)} L_x^{(\alpha)} u_n(x,t) \\ u_0(x,t) = \frac{x^{2\alpha}}{\Gamma(1+2\alpha)} \end{cases} \quad (40)$$

Then, the solution can be expressed as

$$u(x,t) = \frac{x^{2\alpha}}{\Gamma(1+2\alpha)} + \frac{x^\alpha}{\Gamma(1+\alpha)} \frac{t^\alpha}{\Gamma(1+\alpha)} + a^{2\alpha} \frac{t^{2\alpha}}{\Gamma(1+2\alpha)}. \quad (41)$$

The examples of using Adomian decomposition method in solving the integral and differential equations and in heat transfer problems can be seen in [6, 7] and [9]. For examples presented there, one can easily implement local fractional calculus when fractal boundary value problems are considered.

#### 4. Conclusions

Presented integral and differential equations play a very important role in heat conduction problems. Some initial boundary problems for some differential partial equations in physics are reducible to the above integral equation. The development of the investigation for dynamics of the complex systems requires new methods and techniques to be developed. On the other hand many of the classical methods were generalized within the local fractional calculus environment but still the hidden behaviors cannot be revealed properly. In the work, we investigated the fractal initial boundary value problems for local fractional equations with local fractional operators, which are set up on fractals. For these reasons, based on the local fractional operators, the local fractional decomposition method to solve local fractional equations has been applied. The method focuses especially on the approximation methodology for processing local fractional equations. The methodology has been exemplified by four illustrative problems demonstrating the accuracy and the reliabilities of the local fractional decomposition method.

#### Acknowledgement

This research was partially sponsored by the research grant of the Serbian Ministry of Education, Science and Technological Development under the numbers OI 174001, III41006 and TI 35006.

#### Nomenclature

t	time [s]
x	space co-ordinate, [m]
${}_a I_b^{(\alpha)}$	local fractional integral
$L_x^{(\alpha)}$	local fractional order differential operator
$E_\alpha$	Mittag-Leffler function defined on fractal set

## Greek symbols

- $\alpha$  fractal dimension  
 $\Gamma$  gamma function

## References

- [1] Majumder, M., *et al.*, Nanoscale Hydrodynamics – Enhanced Flow in Carbon Nanotubes, *Nature*, 438 (2005), pp. 44, doi: 10.1038/438044a, 7064
- [2] Xuan, Y., Roetzel, W., Conceptions for Heat Transfer Correlation of Nanofluids, *Int. J. Heat Mass Transfer*, 43 (2000), 19, pp. 3701-3707
- [3] Ramšak, M., Leopold, Š., Heat diffusion in fractal geometry cooling surface, *Thermal Science*, 16 (2012), 4, pp. 955-968
- [4] Yang, X. J., Baleanu, D., Fractal heat conduction problem solved by local fractional variation iteration method, *Thermal Science*, 17 (2013), 2, pp. 625-628
- [5] Adomian, G., A review of the decomposition method and some recent results for nonlinear equation, *Mathematical and Computer Modelling*, 13 (1990), 7, pp. 17-43
- [6] Adomian, G., *Solving Frontier Problems of Physics: The Decomposition Method*, Kluwer, Dordrecht, Netherland, 1994
- [7] Wazwaz, A. M., A reliable technique for solving the weakly singular second kind Volterra-type integral equations, *Applied mathematics and computation*, 80 (1996), pp. 287-299
- [8] Luo, X. G., Wu, Q. B., Zhang, B. Q., Revisit on partial solutions in the Adomian decomposition method: Solving heat and wave equations, *Journal of Mathematical Analysis and Applications*, 321 (2006), 1, pp. 353-363
- [9] Nadeem, S., Akbar, N., Effects of heat transfer on the peristaltic transport of MHD Newtonian fluid with variable viscosity: Application of Adomian decomposition method, *Communications in Nonlinear Science and Numerical Simulation*, 14 (2009), 11, pp. 3844-3855
- [10] He, J. H., Variational iteration method-a kind of nonlinear analytical technique: some examples, *International Journal of Non-Linear Mechanics*, 34 (1999), pp. 708-799
- [11] He, J. H., A variational iteration approach to nonlinear problems and its applications, *Mechanical Applications*, 20 (1998), 1, pp. 30-31
- [12] He, J. H., Asymptotic methods for solitary solutions and compactons, *Abstract and Applied Analysis*, 2012 (2012), Article ID 916793, pp. 1-130
- [13] Khan, Y., Mohyud-Din, S. T., Coupling of He's polynomials and Laplace transformation for MHD viscous flow over a stretching sheet, *International Journal of Nonlinear Sciences and Numerical Simulation*, 11 (2010), 12, pp. 1103–1107
- [14] Hristov, J., Heat-Balance integral to fractional (Half-Time) heat diffusion sub-model, *Thermal Science*, 14 (2010), 2, pp. 291-316
- [15] Hristov, J., Approximate solutions to fractional sub-diffusion equations: The heat-balance integral method, *The European Physical Journal-Special Topics*, 193 (2011), pp. 229–243
- [16] Li, Z. B., He, J. H., Fractional complex transform for fractional differential equations, *Mathematical and Computational Applications*, 15 (2010), 5, pp. 970-973
- [17] He, J. H., Elagan S. K., Li, Z. B., Geometrical explanation of the fractional complex transform and derivative chain rule for fractional calculus, *Physics Letters A*, 376 (2012), 4, pp. 257-259

- [18] Jafari H., Seifi, S., Homotopy analysis method for solving linear and nonlinear fractional diffusion-wave equation, *Communications in Nonlinear Science and Numerical Simulation*, 14 (2009), 5, pp. 2006-2012
- [19] Zhang S., Zhang, H. Q., Fractional sub-equation method and its applications to nonlinear fractional PDEs, *Physics Letters A*, 375 (2011), 7, pp. 1069-1073
- [20] Wu, G. C., Lee, E. W. M., Fractional variational iteration method and its application, *Physics Letters A*, 374 (2010), 25, pp. 2506-2509
- [21] Khan, Y., Faraz, N., Yildirim, A., Wu, Q. B., Fractional variational iteration method for fractional initial-boundary value problems arising in the application of nonlinear science, *Computers and Mathematics with Applications*, 62 (2011), 5, pp. 2273-2278
- [22] He, J. H., A short remark on fractional variational iteration method, *Physics Letters A*, 375 (2011), 38, pp. 3362-3364
- [23] Baleanu, D., Diethelm, K., Scalas, E., Trujillo, J. J., *Fractional calculus models and numerical methods*, Series on Complexity, Nonlinearity and Chaos, World Scientific, Boston, USA, 2012
- [24] Kilbas, A. A., Srivastava, H. M., Trujillo, J. J., *Theory and Applications of Fractional Differential Equations*, Elsevier, Amsterdam, Netherland, 2006
- [25] Sabatier, J., Agrawal, O. P., Tenreiro Machado J.A., *Advances in Fractional Calculus: Theoretical Developments and Applications in Physics and Engineering*, Springer, New York, USA, 2007
- [26] Podlubny, I., *Fractional Differential Equations*, Academic Press, New York, USA, 1999
- [27] Yang, X. J., *Local Fractional Functional Analysis and Its Applications*, Asian Academic Publisher Limited, Hong Kong, China , 2011
- [28] Yang, X. J., *Advanced Local Fractional Calculus and Its Applications*, World Science Publisher, New York, USA, 2012.
- [29] Yang, X. J., Local fractional Kernel transform in fractal space and its applications, *Advances in Computational Mathematics and its Applications*, 1 (2012), 2, pp. 86-93
- [30] Yang, X. J., Local fractional integral equations and their applications, *Advances in Computer Science and its Applications*, 1 (2012), 4, pp. 234-239
- [31] Yang, X. J., Zhang, F. R., Local fractional variational iteration method and its algorithms, *Advances in Computational Mathematics and its Applications*, 1 (2012), 3, pp. 139-145
- [32] Yang, X. J., Zhang, Y. D., A new Adomian decomposition procedure scheme for solving local fractional Volterra integral equation, *Advances in Information Technology and Management*, 1 (2012), 4, pp. 158-161
- [33] Yang, X. J., Baleanu, D, Zhong, W. P., Approximation solutions for diffusion equation on Cantor time-space, *Proceeding of the Romanian Academy, Series A*, 14 (2013), 2, pp. 127–133
- [34] Hu, M. S., Agarwal, R. P., Yang, X. J., Local fractional Fourier series with application to wave equation in fractal vibrating string, *Abstract and Applied Analysis*, 2012 (2012), Article ID 567401, pp. 1-15
- [35] Liao, M. K., Yang, X. J., Yan, Q., A new viewpoint to Fourier analysis in fractal space, in: *Advances in Applied Mathematics and Approximation Theory* (Ed. G. A. Anastassiou, O. Duman) Springer, New York, 2013, pp. 399-411

- [36] Zhong, W. P., Gao, F., Shen, X. M., Applications of Yang-Fourier transform to local fractional equations with local fractional derivative and local fractional integral, *Advanced Materials Research*, 461 (2012), pp. 306-310
- [37] Guo, Y., Local fractional Z transform in fractal space, *Advances in Digital Multimedia*, 1 (2012), 2, pp. 96-102.
- [38] Hu, M. S., Baleanu, D., Yang, X. J., One-phase problems for discontinuous heat transfer in fractal media, *Mathematical Problems in Engineering*, 2013 (2013), Article ID 358473, pp. 1-3
- [39] Yang, A. M., Yang, X. J., Li, Z. B., Local fractional series expansion method for solving wave and diffusion equations on Cantor set, *Abstract and Applied Analysis*, 2013 (2013), Article ID 351057, pp. 1–5
- [40] Yang, X. J., Baleanu, D., Tenreiro Machado J. A., Mathematical aspects of the Heisenberg uncertainty principle within local fractional Fourier analysis, *Boundary Value Problems*, 2013 (2013), 131, pp. 1–16
- [41] Yang, Y. J., Baleanu, D, Yang, X. J., Analysis of wave equations by local fractional Fourier series method, *Advances in Mathematical Physics*, 2013 (2013), Article ID 632309, pp. 1–7