

## *Research Article*

# **Common Fixed Point Theorems for Four Mappings on Cone Metric Type Space**

**Aleksandar S. Cvetković,<sup>1</sup> Marija P. Stanić,<sup>2</sup>  
Sladjana Dimitrijević,<sup>2</sup> and Suzana Simić<sup>2</sup>**

<sup>1</sup> *Department of Mathematics, Faculty of Mechanical Engineering, University of Belgrade, Kraljice Marije 16, 11120 Belgrade, Serbia*

<sup>2</sup> *Department of Mathematics and Informatics, Faculty of Science, University of Kragujevac, Radoja Domanovića 12, 34000 Kragujevac, Serbia*

Correspondence should be addressed to Aleksandar S. Cvetković, [acvetkovic@mas.bg.ac.rs](mailto:acvetkovic@mas.bg.ac.rs)

Received 9 December 2010; Revised 26 January 2011; Accepted 3 February 2011

Academic Editor: Fabio Zanolin

Copyright © 2011 Aleksandar S. Cvetković et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In this paper we consider the so called a cone metric type space, which is a generalization of a cone metric space. We prove some common fixed point theorems for four mappings in those spaces. Obtained results extend and generalize well-known comparable results in the literature. All results are proved in the settings of a solid cone, without the assumption of continuity of mappings.

## **1. Introduction**

Replacing the real numbers, as the codomain of a metric, by an ordered Banach space we obtain a generalization of metric space. Such a generalized space, called a cone metric space, was introduced by Huang and Zhang in [1]. They described the convergence in cone metric space, introduced their completeness, and proved some fixed point theorems for contractive mappings on cone metric space. Cones and ordered normed spaces have some applications in optimization theory (see [2]). The initial work of Huang and Zhang [1] inspired many authors to prove fixed point theorems, as well as common fixed point theorems for two or more mappings on cone metric space, for example, [3–14].

In this paper we consider the so-called a cone metric type space, which is a generalization of a cone metric space and prove some common fixed point theorems for four mappings in those spaces. Obtained results are generalization of theorems proved in [13]. For some special choices of mappings we obtain theorems which generalize results from [1, 8, 15].

All results are proved in the settings of a solid cone, without the assumption of continuity of mappings.

The paper is organized as follows. In Section 2 we repeat some definitions and well-known results which will be needed in the sequel. In Section 3 we prove common fixed point theorems. Also, we presented some corollaries which show that our results are generalization of some existing results in the literature.

## 2. Definitions and Notation

Let  $E$  be a real Banach space and  $P$  a subset of  $E$ . By  $\theta$  we denote zero element of  $E$  and by  $\text{int } P$  the interior of  $P$ . The subset  $P$  is called a *cone* if and only if

- (i)  $P$  is closed, nonempty and  $P \neq \{\theta\}$ ;
- (ii)  $a, b \in \mathbb{R}$ ,  $a, b \geq 0$ , and  $x, y \in P$  imply  $ax + by \in P$ ;
- (iii)  $P \cap (-P) = \{\theta\}$ .

For a given cone  $P$ , a partial ordering  $\leq$  with respect to  $P$  is introduced in the following way:  $x \leq y$  if and only if  $y - x \in P$ . One writes  $x < y$  to indicate that  $x \leq y$ , but  $x \neq y$ . If  $y - x \in \text{int } P$ , one writes  $x \ll y$ .

If  $\text{int } P \neq \emptyset$ , the cone  $P$  is called *solid*.

In the sequel we always suppose that  $E$  is a real Banach space,  $P$  is a solid cone in  $E$ , and  $\leq$  is partial ordering with respect to  $P$ .

Analogously with definition of metric type space, given in [16], we consider cone metric type space.

*Definition 2.1.* Let  $X$  be a nonempty set and  $E$  a real Banach space with cone  $P$ . A vector-valued function  $d : X \times X \rightarrow E$  is said to be a cone metric type function on  $X$  with constant  $K \geq 1$  if the following conditions are satisfied:

- ( $d_1$ )  $\theta \leq d(x, y)$  for all  $x, y \in X$  and  $d(x, y) = \theta$  if and only if  $x = y$ ;
- ( $d_2$ )  $d(x, y) = d(y, x)$  for all  $x, y \in X$ ;
- ( $d_3$ )  $d(x, y) \leq K(d(x, z) + d(z, y))$  for all  $x, y, z \in X$ .

The pair  $(X, d)$  is called a cone metric type space (in brief CMTS).

*Remark 2.2.* For  $K = 1$  in Definition 2.1 we obtain a cone metric space introduced in [1].

*Definition 2.3.* Let  $(X, d)$  be a CMTS and  $\{x_n\}$  a sequence in  $X$ .

- ( $c_1$ )  $\{x_n\}$  converges to  $x \in X$  if for every  $c \in E$  with  $\theta \ll c$  there exists  $n_0 \in \mathbb{N}$  such that  $d(x_n, x) \ll c$  for all  $n > n_0$ . We write  $\lim_{n \rightarrow \infty} x_n = x$ , or  $x_n \rightarrow x, n \rightarrow \infty$ .
- ( $c_2$ ) If for every  $c \in E$  with  $\theta \ll c$  there exists  $n_0 \in \mathbb{N}$  such that  $d(x_n, x_m) \ll c$  for all  $n, m > n_0$ , then  $\{x_n\}$  is called a Cauchy sequence in  $X$ .

If every Cauchy sequence is convergent in  $X$ , then  $X$  is called a complete CMTS.

*Example 2.4.* Let  $B = \{e_i \mid i = 1, \dots, n\}$  be orthonormal basis of  $\mathbb{R}^n$  with inner product  $(\cdot, \cdot)$ . Let  $p > 0$ , and define

$$X_p = \left\{ [x] \mid x : [0, 1] \longrightarrow \mathbb{R}^n, \int_0^1 |(x(t), e_k)|^p dt \in \mathbb{R}, k = 1, \dots, n \right\}, \quad (2.1)$$

where  $[x]$  represents class of element  $x$  with respect to equivalence relation of functions equal almost everywhere. We choose  $E = \mathbb{R}^n$  and

$$P_B = \{y \in \mathbb{R}^n \mid (y, e_i) \geq 0, i = 1, \dots, n\}. \quad (2.2)$$

We show that  $P_B$  is a solid cone. Let  $y_k \in P_B, k \in \mathbb{N}$ , with property  $\lim_{k \rightarrow +\infty} y_k = y$ . Since scalar product is continuous, we get  $\lim_{k \rightarrow +\infty} (y_k, e_i) = (\lim_{k \rightarrow +\infty} y_k, e_i) = (y, e_i), i = 1, \dots, n$ . Clearly, it must be  $(y, e_i) \geq 0, i = 1, \dots, n$ , and, hence,  $y \in P_B$ , that is,  $P_B$  is closed. It is obvious that  $\theta \neq e_1 \in P_B \neq \{\theta\}$ , and for  $a, b \geq 0$ , and all  $z, y \in P_B$ , we have  $(az + by, e_i) = a(z, e_i) + b(y, e_i) \geq 0, i = 1, \dots, n$ . Finally, if  $z \in P_B \cap (-P_B)$  we have  $(z, e_i) \geq 0$  and  $(-z, e_i) \geq 0, i = 1, \dots, n$ , and it follows that  $(z, e_i) = 0, i = 1, \dots, n$ , and, since  $B$  is complete, we get  $z = 0$ . Let us choose  $z = \sum_{i=1}^n e_i$ . It is obvious that  $z \in \text{int } P_B$ , since if not, for every  $\varepsilon > 0$  there exists  $y \notin P_B$  such that  $|1 - (y, e_i)| \leq (\sum_{i=1}^n |1 - (y, e_i)|^2)^{1/2} = \|z - y\| < \varepsilon$ . If we choose  $\varepsilon = 1/4$ , we conclude that it must be  $(y, e_i) > 1 - 1/4 > 0$ , hence  $y \in P_B$ , which is contradiction.

Finally, define  $d : X_p \times X_p \rightarrow P_B$  by

$$d(f, g) = \sum_{i=1}^n e_i \int_0^1 |((f - g)(t), e_i)|^p dt, \quad f, g \in X_p. \quad (2.3)$$

Then it is obvious that  $(X_p, d)$  is CMTS with  $K = 2^{p-1}$ . Let  $f, g, h$  be functions such that  $(f, e_1) = 1, (g, e_1) = -2, (h, e_1) = 0$ , and  $(f, e_i) = (g, e_i) = (h, e_i) = 0, i = 2, \dots, n$ , with  $p = 2$  give  $d(f, g) = 9e_1, d(f, h) = e_1$ , and  $d(h, g) = 4e_1$ , which proves  $5e_1 = d(f, h) + d(h, g) \leq d(f, g) = 9e_1$ , but  $9e_1 = d(f, g) \leq 2(d(f, h) + d(h, g)) = 10e_1$ .

The following properties are well known in the case of a cone metric space, and it is easy to see that they hold also in the case of a CMTS.

**Lemma 2.5.** *Let  $(X, d)$  be a CMTS over-ordered real Banach space  $E$  with a cone  $P$ . The following properties hold ( $a, b, c \in E$ ).*

- (p<sub>1</sub>) *If  $a \leq b$  and  $b \ll c$ , then  $a \ll c$ .*
- (p<sub>2</sub>) *If  $\theta \leq a \ll c$  for all  $c \in \text{int } P$ , then  $a = \theta$ .*
- (p<sub>3</sub>) *If  $a \leq \lambda a$ , where  $a \in P$  and  $0 \leq \lambda < 1$ , then  $a = \theta$ .*
- (p<sub>4</sub>) *Let  $x_n \rightarrow \theta$  in  $E$  and let  $\theta \ll c$ . Then there exists positive integer  $n_0$  such that  $x_n \ll c$  for each  $n > n_0$ .*

*Definition 2.6* (see [17]). Let  $F, G : X \rightarrow X$  be mappings of a set  $X$ . If  $y = Fx = Gx$  for some  $x \in X$ , then  $x$  is called a coincidence point of  $F$  and  $G$ , and  $y$  is called a point of coincidence of  $F$  and  $G$ .

*Definition 2.7* (see [17]). Let  $F$  and  $G$  be self-mappings of set  $X$  and  $\mathcal{C}(F, G) = \{x \in X : Fx = Gx\}$ . The pair  $\{F, G\}$  is called weakly compatible if mappings  $F$  and  $G$  commute at all their coincidence points, that is, if  $FGx = GFx$  for all  $x \in \mathcal{C}(F, G)$ .

**Lemma 2.8** (see [5]). *Let  $F$  and  $G$  be weakly compatible self-mappings of a set  $X$ . If  $F$  and  $G$  have a unique point of coincidence  $y = Fx = Gx$ , then  $y$  is the unique common fixed point of  $F$  and  $G$ .*

### 3. Main Results

**Theorem 3.1.** *Let  $(X, d)$  be a CMTS with constant  $1 \leq K \leq 2$  and  $P$  a solid cone. Suppose that self-mappings  $F, G, S, T : X \rightarrow X$  are such that  $SX \subset GX$ ,  $TX \subset FX$  and that for some constant  $\lambda \in (0, 1/K)$  for all  $x, y \in X$  there exists*

$$u(x, y) \in \left\{ Kd(Fx, Gy), Kd(Fx, Sx), Kd(Gy, Ty), K \frac{d(Fx, Ty) + d(Gy, Sx)}{2} \right\}, \quad (3.1)$$

such that the following inequality

$$d(Sx, Ty) \leq \frac{\lambda}{K} u(x, y), \quad (3.2)$$

holds. If one of  $SX$ ,  $TX$ ,  $FX$ , or  $GX$  is complete subspace of  $X$ , then  $\{S, F\}$  and  $\{T, G\}$  have a unique point of coincidence in  $X$ . Moreover, if  $\{S, F\}$  and  $\{T, G\}$  are weakly compatible pairs, then  $F, G, S$ , and  $T$  have a unique common fixed point.

*Proof.* Let us choose  $x_0 \in X$  arbitrary. Since  $SX \subset GX$ , there exists  $x_1 \in X$  such that  $Gx_1 = Sx_0 = z_0$ . Since  $TX \subset FX$ , there exists  $x_2 \in X$  such that  $Fx_2 = Tx_1 = z_1$ . We continue in this manner. In general,  $x_{2n+1} \in X$  is chosen such that  $Gx_{2n+1} = Sx_{2n} = z_{2n}$ , and  $x_{2n+2} \in X$  is chosen such that  $Fx_{2n+2} = Tx_{2n+1} = z_{2n+1}$ .

First we prove that

$$d(z_n, z_{n+1}) \leq \alpha d(z_{n-1}, z_n), \quad n \geq 1, \quad (3.3)$$

where  $\alpha = \max\{\lambda, \lambda K / (2 - \lambda K)\}$ , which will lead us to the conclusion that  $\{z_n\}$  is a Cauchy sequence, since  $\alpha \in (0, 1)$  (it is easy to see that  $0 < \lambda K / (2 - \lambda K) < 1$ ). To prove this, it is necessary to consider the cases of an odd integer  $n$  and of an even  $n$ .

For  $n = 2\ell + 1$ ,  $\ell \in \mathbb{N}_0$ , we have  $d(z_{2\ell+1}, z_{2\ell+2}) = d(Sx_{2\ell+2}, Tx_{2\ell+1})$ , and from (3.2) there exists

$$\begin{aligned} u(x_{2\ell+2}, x_{2\ell+1}) &\in \left\{ Kd(Fx_{2\ell+2}, Gx_{2\ell+1}), Kd(Fx_{2\ell+2}, Sx_{2\ell+2}), \right. \\ &\quad \left. Kd(Gx_{2\ell+1}, Tx_{2\ell+1}), K \frac{d(Fx_{2\ell+2}, Tx_{2\ell+1}) + d(Gx_{2\ell+1}, Sx_{2\ell+2})}{2} \right\} \quad (3.4) \\ &= \left\{ Kd(z_{2\ell+1}, z_{2\ell}), Kd(z_{2\ell+1}, z_{2\ell+2}), \frac{Kd(z_{2\ell}, z_{2\ell+2})}{2} \right\}, \end{aligned}$$

such that  $d(z_{2\ell+1}, z_{2\ell+2}) \leq (\lambda/K)u(x_{2\ell+2}, x_{2\ell+1})$ . Thus we have the following three cases:

- (i)  $d(z_{2\ell+1}, z_{2\ell+2}) \leq \lambda d(z_{2\ell+1}, z_{2\ell})$ ;
- (ii)  $d(z_{2\ell+1}, z_{2\ell+2}) \leq \lambda d(z_{2\ell+1}, z_{2\ell+2})$ , which, because of property  $(p_3)$ , implies  $d(z_{2\ell+1}, z_{2\ell+2}) = \theta$ ;
- (iii)  $d(z_{2\ell+1}, z_{2\ell+2}) \leq (\lambda/2)d(z_{2\ell}, z_{2\ell+2})$ , that is, by using  $(d_3)$ ,

$$d(z_{2\ell+1}, z_{2\ell+2}) \leq \frac{\lambda K}{2}d(z_{2\ell}, z_{2\ell+1}) + \frac{\lambda K}{2}d(z_{2\ell+1}, z_{2\ell+2}), \quad (3.5)$$

which implies  $d(z_{2\ell+1}, z_{2\ell+2}) \leq (\lambda K/(2 - \lambda K))d(z_{2\ell}, z_{2\ell+1})$ .

Thus, inequality (3.3) holds in this case.

For  $n = 2\ell$ ,  $\ell \in \mathbb{N}_0$ , we have

$$d(z_{2\ell}, z_{2\ell+1}) = d(Sx_{2\ell}, Tx_{2\ell+1}) \leq \frac{\lambda}{K}u(x_{2\ell}, x_{2\ell+1}), \quad (3.6)$$

where

$$\begin{aligned} u(x_{2\ell}, x_{2\ell+1}) &\in \left\{ Kd(Fx_{2\ell}, Gx_{2\ell+1}), Kd(Fx_{2\ell}, Sx_{2\ell}), \right. \\ &\quad \left. Kd(Gx_{2\ell+1}, Tx_{2\ell+1}), K \frac{d(Fx_{2\ell}, T_{2\ell+1}) + d(Gx_{2\ell+1}, Sx_{2\ell})}{2} \right\} \quad (3.7) \\ &= \left\{ Kd(z_{2\ell-1}, z_{2\ell}), Kd(z_{2\ell}, z_{2\ell+1}), \frac{Kd(z_{2\ell-1}, z_{2\ell+1})}{2} \right\}. \end{aligned}$$

Thus we have the following three cases:

- (i)  $d(z_{2\ell}, z_{2\ell+1}) \leq \lambda d(z_{2\ell-1}, z_{2\ell})$ ;
- (ii)  $d(z_{2\ell}, z_{2\ell+1}) \leq \lambda d(z_{2\ell}, z_{2\ell+1})$ , which implies  $d(z_{2\ell}, z_{2\ell+1}) = \theta$ ;
- (iii)  $d(z_{2\ell}, z_{2\ell+1}) \leq (\lambda/2)d(z_{2\ell-1}, z_{2\ell+1}) \leq (\lambda K/2)d(z_{2\ell-1}, z_{2\ell}) + (\lambda K/2)d(z_{2\ell}, z_{2\ell+1})$ , which implies  $d(z_{2\ell}, z_{2\ell+1}) \leq (\lambda K/(2 - \lambda K))d(z_{2\ell}, z_{2\ell-1})$ .

So, inequality (3.3) is satisfied in this case, too.

Therefore, (3.3) is satisfied for all  $n \in \mathbb{N}_0$ , and by iterating we get

$$d(z_n, z_{n+1}) \leq \alpha^n d(z_0, z_1). \quad (3.8)$$

Since  $K \geq 1$ , for  $m > n$  we have

$$\begin{aligned} d(z_n, z_m) &\leq Kd(z_n, z_{n+1}) + K^2d(z_{n+1}, z_{n+2}) + \cdots + K^{m-n-1}d(z_{m-1}, z_m) \\ &\leq \left( K\alpha^n + K^2\alpha^{n+1} + \cdots + K^{m-n}\alpha^{m-1} \right) d(z_0, z_1) \quad (3.9) \\ &\leq \frac{K\alpha^n}{1 - K\alpha} d(z_0, z_1) \longrightarrow \theta, \quad \text{as } n \longrightarrow \infty. \end{aligned}$$

Now, by  $(p_4)$  and  $(p_1)$ , it follows that for every  $c \in \text{int } P$  there exists positive integer  $n_0$  such that  $d(z_n, z_m) \ll c$  for every  $m > n > n_0$ , so  $\{z_n\}$  is a Cauchy sequence.

Let us suppose that  $SX$  is complete subspace of  $X$ . Completeness of  $SX$  implies existence of  $z \in SX$  such that  $\lim_{n \rightarrow \infty} z_{2n} = \lim_{n \rightarrow \infty} Sx_{2n} = z$ . Then, we have

$$\lim_{n \rightarrow \infty} Gx_{2n+1} = \lim_{n \rightarrow \infty} Sx_{2n} = \lim_{n \rightarrow \infty} Fx_{2n} = \lim_{n \rightarrow \infty} Tx_{2n+1} = z, \quad (3.10)$$

that is, for any  $\theta \ll c$ , for sufficiently large  $n$  we have  $d(z_n, z) \ll c$ . Since  $z \in SX \subset GX$ , there exists  $y \in X$  such that  $z = Gy$ . Let us prove that  $z = Ty$ . From  $(d_3)$  and (3.2), we have

$$d(Ty, z) \leq Kd(Ty, Sx_{2n}) + Kd(Sx_{2n}, z) \leq \lambda u(x_{2n}, y) + Kd(z_{2n}, z), \quad (3.11)$$

where

$$\begin{aligned} u(x_{2n}, y) &\in \left\{ Kd(Fx_{2n}, Gy), Kd(Fx_{2n}, Sx_{2n}), Kd(Gy, Ty), K \frac{d(Fx_{2n}, Ty) + d(Gy, Sx_{2n})}{2} \right\} \\ &= \left\{ Kd(z_{2n-1}, z), Kd(z_{2n-1}, z_{2n}), Kd(z, Ty), K \frac{d(z_{2n-1}, Ty) + d(z, z_{2n})}{2} \right\}. \end{aligned} \quad (3.12)$$

Therefore we have the following four cases:

- (i)  $d(Ty, z) \leq K\lambda d(z_{2n-1}, z) + Kd(z_{2n}, z) \ll K\lambda \cdot c / (2K\lambda) + K \cdot c / (2K) = c$ , as  $n \rightarrow \infty$ ;
- (ii)  $d(Ty, z) \leq K\lambda d(z_{2n-1}, z_{2n}) + Kd(z_{2n}, z) \ll K\lambda \cdot c / (2K\lambda) + K \cdot c / (2K) = c$ , as  $n \rightarrow \infty$ ;
- (iii)  $d(Ty, z) \leq K\lambda d(z, Ty) + Kd(z_{2n}, z)$ , that is,

$$d(Ty, z) \leq \frac{K}{1-K\lambda} d(z_{2n}, z) \ll \frac{K}{1-K\lambda} \cdot \frac{1-K\lambda}{K} \cdot c = c, \quad \text{as } n \rightarrow \infty; \quad (3.13)$$

- (iv)  $d(Ty, z) \leq (K\lambda/2)(d(z_{2n-1}, Ty) + d(z, z_{2n})) + Kd(z_{2n}, z)$ , that is, because of  $(d_3)$ ,

$$d(Ty, z) \leq \frac{K\lambda}{2} (Kd(z_{2n-1}, z) + Kd(z, Ty) + d(z, z_{2n})) + Kd(z_{2n}, z), \quad (3.14)$$

which implies

$$\begin{aligned} d(Ty, z) &\leq \frac{1}{1-K^2\lambda/2} \left[ \frac{K^2\lambda}{2} d(z_{2n-1}, z) + \left( \frac{K\lambda}{2} + K \right) d(z_{2n}, z) \right] \\ &\ll \frac{K^2\lambda}{2-K^2\lambda} \frac{2-K^2\lambda}{K^2\lambda} \frac{c}{2} + \frac{K(\lambda+2)}{2-K^2\lambda} \frac{2-K^2\lambda}{K(\lambda+2)} \frac{c}{2} = c, \quad \text{as } n \rightarrow \infty, \end{aligned} \quad (3.15)$$

since from  $1 \leq K \leq 2$  and  $\lambda \in (0, 1/K)$  we have  $\lambda < 1/K \leq 2/K^2$ , and therefore  $1 - K^2\lambda/2 > 0$ .

Therefore,  $d(Ty, z) \ll c$  for each  $c \in \text{int } P$ . So, by  $(p_2)$  we have  $d(Ty, z) = \theta$ , that is,  $Ty = Gy = z$ ,  $y$  is a coincidence point, and  $z$  is a point of coincidence of  $T$  and  $G$ .

Since  $TX \subset FX$ , there exists  $v \in X$  such that  $z = Fv$ . Let us prove that  $Sv = z$ . From  $(d_3)$  and (3.2), we have

$$d(Sv, z) \leq Kd(Sv, Tx_{2n+1}) + Kd(Tx_{2n+1}, z) \leq \lambda u(v, x_{2n+1}) + Kd(z_{2n+1}, z), \quad (3.16)$$

where

$$\begin{aligned} u(v, x_{2n+1}) & \in \left\{ Kd(Fv, Gx_{2n+1}), Kd(Fv, Sv), Kd(Gx_{2n+1}, Tx_{2n+1}), K \frac{d(Fv, Tx_{2n+1}) + d(Gx_{2n+1}, Sv)}{2} \right\} \\ & = \left\{ Kd(z, z_{2n}), Kd(z, Sv), Kd(z_{2n}, z_{2n+1}), K \frac{d(z, z_{2n+1}) + d(z_{2n}, Sv)}{2} \right\}. \end{aligned} \quad (3.17)$$

Therefore we have the following four cases:

- (i)  $d(Sv, z) \leq K\lambda d(z, z_{2n}) + Kd(z_{2n+1}, z)$ ;
- (ii)  $d(Sv, z) \leq K\lambda d(z, Sv) + Kd(z_{2n+1}, z)$ ;
- (iii)  $d(Sv, z) \leq K\lambda d(z_{2n}, z_{2n+1}) + Kd(z_{2n+1}, z)$ ;
- (iv)  $d(Sv, z) \leq (K\lambda/2)(d(z, z_{2n+1}) + d(z_{2n}, Sv)) + Kd(z_{2n+1}, z)$ .

By the same arguments as above, we conclude that  $d(Sv, z) = \theta$ , that is,  $Sv = Fv = z$ . So,  $z$  is a point of coincidence of  $S$  and  $F$ , too.

Now we prove that  $z$  is unique point of coincidence of pairs  $\{S, F\}$  and  $\{T, G\}$ . Suppose that there exists  $z^*$  which is also a point of coincidence of these four mappings, that is,  $Fv^* = Gy^* = Sv^* = Ty^* = z^*$ . From (3.2),

$$d(z, z^*) = d(Sv, Ty^*) \leq \frac{\lambda}{K} u(v, y^*), \quad (3.18)$$

where

$$\begin{aligned} u(v, y^*) & \in \left\{ Kd(Fv, Gy^*), Kd(Fv, Sv), d(Gy^*, Ty^*), K \frac{d(Fv, Ty^*) + d(Gy^*, Sv)}{2} \right\} \\ & = \{Kd(z, z^*), \theta\}. \end{aligned} \quad (3.19)$$

Using  $(p_3)$  we get  $d(z, z^*) = \theta$ , that is,  $z = z^*$ . Therefore,  $z$  is the unique point of coincidence of pairs  $\{S, F\}$  and  $\{T, G\}$ . If these pairs are weakly compatible, then  $z$  is the unique common fixed point of  $S, F, T$ , and  $G$ , by Lemma 2.8.

Similarly, we can prove the statement in the cases when  $FX, GX$ , or  $TX$  is complete.  $\square$

We give one simple, but illustrative, example.

*Example 3.2.* Let  $X = \mathbb{R}$ ,  $E = \mathbb{R}$ , and  $P = [0, +\infty)$ . Let us define  $d(x, y) = |x - y|^2$  for all  $x, y \in X$ . Then  $(X, d)$  is a CMTS, but it is not a cone metric space since the triangle inequality is not satisfied. Starting with Minkowski inequality (see [18]) for  $p = 2$ , by using the inequality of arithmetic and geometric means, we get

$$|x - z|^2 \leq |x - y|^2 + |y - z|^2 + 2|x - y||x - z| \leq 2(|x - y|^2 + |y - z|^2). \quad (3.20)$$

Here,  $K = 2$ .

Let us define four mappings  $S, F, T, G : X \rightarrow X$  as follows:

$$Sx = M(ax + b), \quad Fx = ax + b, \quad Tx = M(cx + d), \quad Gx = cx + d, \quad (3.21)$$

where  $x \in X$ ,  $a \neq 0$ ,  $c \neq 0$ , and  $M < 1/\sqrt{2}$ . Since  $SX = FX = TX = GX = X$  we have trivially  $SX \subset GX$  and  $TX \subset FX$ . Also,  $X$  is a complete space. Further,  $d(Sx, Ty) = |M(ax + b) - M(cy + d)|^2 = M^2d(Fx, Gy)$ , that is, there exists  $\lambda = M^2 < 1/2 = 1/K$  such that (3.2) is satisfied.

According to Theorem 3.1,  $\{S, F\}$  and  $\{T, G\}$  have a unique point of coincidence in  $X$ , that is, there exists unique  $z \in X$  and there exist  $x, y \in X$  such that  $z = Sx = Fx = Ty = Gy$ . It is easy to see that  $x = -b/a$ ,  $y = -d/c$ , and  $z = 0$ .

If  $\{S, F\}$  is weakly compatible pair, we have  $SFx = FSx$ , which implies  $Mb = b$ , that is,  $b = 0$ . Similarly, if  $\{T, G\}$  is weakly compatible pair, we have  $TGy = GTy$ , which implies  $Md = d$ , that is,  $d = 0$ . Then  $x = y = 0$ , and  $z = 0$  is the unique common fixed point of these four mappings.

The following two theorems can be proved in the same way as Theorem 3.1, so we omit the proofs.

**Theorem 3.3.** *Let  $(X, d)$  be a CMTS with constant  $K \geq 2$  and  $P$  a solid cone. Suppose that self-mappings  $F, G, S, T : X \rightarrow X$  are such that  $SX \subset GX$ ,  $TX \subset FX$  and that for some constant  $\lambda \in (0, 2/K^2)$  for all  $x, y \in X$  there exists*

$$u(x, y) \in \left\{ Kd(Fx, Gy), Kd(Fx, Sx), Kd(Gy, Ty), K \frac{d(Fx, Ty) + d(Gy, Sx)}{2} \right\}, \quad (3.22)$$

such that the following inequality

$$d(Sx, Ty) \leq \frac{\lambda}{K} u(x, y), \quad (3.23)$$

holds. If one of  $SX$ ,  $TX$ ,  $FX$ , or  $GX$  is complete subspace of  $X$ , then  $\{S, F\}$  and  $\{T, G\}$  have a unique point of coincidence in  $X$ . Moreover, if  $\{S, F\}$  and  $\{T, G\}$  are weakly compatible pairs, then  $F, G, S$ , and  $T$  have a unique common fixed point.



**Theorem 3.4.** Let  $(X, d)$  be a CMTS with constant  $K \geq 1$  and  $P$  a solid cone. Suppose that self-mappings  $F, G, S, T : X \rightarrow X$  are such that  $SX \subset GX$ ,  $TX \subset FX$  and that for some constant  $\lambda \in (0, 1/K)$  for all  $x, y \in X$  there exists

$$u(x, y) \in \left\{ Kd(Fx, Gy), Kd(Fx, Sx), Kd(Gy, Ty), \frac{d(Fx, Ty) + d(Gy, Sx)}{2} \right\}, \quad (3.24)$$

such that the following inequality

$$d(Sx, Ty) \leq \frac{\lambda}{K} u(x, y), \quad (3.25)$$

holds. If one of  $SX$ ,  $TX$ ,  $FX$ , or  $GX$  is complete subspace of  $X$ , then  $\{S, F\}$  and  $\{T, G\}$  have a unique point of coincidence in  $X$ . Moreover, if  $\{S, F\}$  and  $\{T, G\}$  are weakly compatible pairs, then  $F, G, S$ , and  $T$  have a unique common fixed point.

Theorems 3.1 and 3.4 are generalizations of [13, Theorem 2.2]. As a matter of fact, for  $K = 1$ , from Theorems 3.1 and 3.4, we get [13, Theorem 2.2].

If we choose  $T = S$  and  $G = F$ , from Theorems 3.1, 3.3, and 3.4 we get the following results for two mappings on CMTS.

**Corollary 3.5.** Let  $(X, d)$  be a CMTS with constant  $1 \leq K \leq 2$  and  $P$  a solid cone. Suppose that self-mappings  $F, S : X \rightarrow X$  are such that  $SX \subset FX$  and that for some constant  $\lambda \in (0, 1/K)$  for all  $x, y \in X$  there exists

$$u(x, y) \in \left\{ Kd(Fx, Fy), Kd(Fx, Sx), Kd(Fy, Sy), K \frac{d(Fx, Sy) + d(Fy, Sx)}{2} \right\}, \quad (3.26)$$

such that the following inequality

$$d(Sx, Sy) \leq \frac{\lambda}{K} u(x, y), \quad (3.27)$$

holds. If  $FX$  or  $SX$  is complete subspace of  $X$ , then  $F$  and  $S$  have a unique point of coincidence in  $X$ . Moreover, if  $\{F, S\}$  is a weakly compatible pair, then  $F$  and  $S$  have a unique common fixed point.

**Corollary 3.6.** Let  $(X, d)$  be a CMTS with constant  $K \geq 2$  and  $P$  a solid cone. Suppose that self-mappings  $F, S : X \rightarrow X$  are such that  $SX \subset FX$  and that for some constant  $\lambda \in (0, 2/K^2)$  for all  $x, y \in X$  there exists

$$u(x, y) \in \left\{ Kd(Fx, Fy), Kd(Fx, Sx), Kd(Fy, Sy), K \frac{d(Fx, Sy) + d(Fy, Sx)}{2} \right\}, \quad (3.28)$$

such that the following inequality

$$d(Sx, Sy) \leq \frac{\lambda}{K} u(x, y), \quad (3.29)$$

holds. If  $FX$  or  $SX$  is complete subspace of  $X$ , then  $F$  and  $S$  have a unique point of coincidence in  $X$ . Moreover, if  $\{F, S\}$  is a weakly compatible pair, then  $F$  and  $S$  have a unique common fixed point.

**Corollary 3.7.** Let  $(X, d)$  be a CMTS with constant  $K \geq 1$  and  $P$  a solid cone. Suppose that self-mappings  $F, S : X \rightarrow X$  are such that  $SX \subset FX$  and that for some constant  $\lambda \in (0, 1/K)$  for all  $x, y \in X$  there exists

$$u(x, y) \in \left\{ Kd(Fx, Fy), Kd(Fx, Sx), Kd(Fy, Sy), \frac{d(Fx, Sy) + d(Fy, Sx)}{2} \right\}, \quad (3.30)$$

such that the following inequality

$$d(Sx, Sy) \leq \frac{\lambda}{K} u(x, y), \quad (3.31)$$

holds. If  $FX$  or  $SX$  is complete subspace of  $X$ , then  $F$  and  $S$  have a unique point of coincidence in  $X$ . Moreover, if  $\{F, S\}$  is a weakly compatible pair, then  $F$  and  $S$  have a unique common fixed point.

**Theorem 3.8.** Let  $(X, d)$  be a CMTS with constant  $K \geq 1$  and  $P$  a solid cone. Suppose that self-mappings  $F, G, S, T : X \rightarrow X$  are such that  $SX \subset GX$ ,  $TX \subset FX$  and that there exist nonnegative constants  $a_i, i = 1, \dots, 5$ , satisfying

$$a_1 + a_2 + a_3 + 2K \max\{a_4, a_5\} < 1, \quad a_3K + a_4K^2 < 1, \quad a_2K + a_5K^2 < 1, \quad (3.32)$$

such that for all  $x, y \in X$  inequality

$$d(Sx, Ty) \leq a_1d(Fx, Gy) + a_2d(Fx, Sx) + a_3d(Gy, Ty) + a_4d(Fx, Ty) + a_5d(Gy, Sx), \quad (3.33)$$

holds. If one of  $SX, TX, FX$ , or  $GX$  is complete subspace of  $X$ , then  $\{S, F\}$  and  $\{T, G\}$  have a unique point of coincidence in  $X$ . Moreover, if  $\{S, F\}$  and  $\{T, G\}$  are weakly compatible pairs, then  $F, G, S$ , and  $T$  have a unique common fixed point.

*Proof.* We define sequences  $\{x_n\}$  and  $\{z_n\}$  as in the proof of Theorem 3.1. First we prove that

$$d(z_n, z_{n+1}) \leq \alpha d(z_{n-1}, z_n), \quad n \geq 1, \quad (3.34)$$

where

$$\alpha = \max \left\{ \frac{a_1 + a_3 + a_5K}{1 - a_2 - a_5K}, \frac{a_1 + a_2 + a_4K}{1 - a_3 - a_4K} \right\}, \quad (3.35)$$

which implies that  $\{z_n\}$  is a Cauchy sequence, since, because of (3.32), it is easy to check that  $\alpha \in [0, 1)$ . To prove this, it is necessary to consider the cases of an odd and of an even integer  $n$ .

For  $n = 2\ell + 1$ ,  $\ell \in \mathbb{N}_0$ , we have  $d(z_{2\ell+1}, z_{2\ell+2}) = d(Sx_{2\ell+2}, Tx_{2\ell+1})$ , and from (3.33) we have

$$\begin{aligned} d(Sx_{2\ell+2}, Tx_{2\ell+1}) &\leq a_1 d(Fx_{2\ell+2}, Gx_{2\ell+1}) + a_2 d(Fx_{2\ell+2}, Sx_{2\ell+2}) \\ &\quad + a_3 d(Gx_{2\ell+1}, Tx_{2\ell+1}) + a_4 d(Fx_{2\ell+2}, Tx_{2\ell+1}) + a_5 d(Gx_{2\ell+1}, Sx_{2\ell+2}), \end{aligned} \quad (3.36)$$

that is,

$$\begin{aligned} d(z_{2\ell+1}, z_{2\ell+2}) &\leq a_1 d(z_{2\ell+1}, z_{2\ell}) + a_2 d(z_{2\ell+1}, z_{2\ell+2}) + a_3 d(z_{2\ell}, z_{2\ell+1}) \\ &\quad + a_4 d(z_{2\ell+1}, z_{2\ell+1}) + a_5 d(z_{2\ell}, z_{2\ell+2}) \\ &= (a_1 + a_3) d(z_{2\ell}, z_{2\ell+1}) + a_2 d(z_{2\ell+1}, z_{2\ell+2}) + a_5 d(z_{2\ell}, z_{2\ell+2}) \\ &\leq (a_1 + a_3) d(z_{2\ell}, z_{2\ell+1}) + a_2 d(z_{2\ell+1}, z_{2\ell+2}) + a_5 K d(z_{2\ell}, z_{2\ell+1}) \\ &\quad + a_5 K d(z_{2\ell+1}, z_{2\ell+2}) \\ &= (a_1 + a_3 + a_5 K) d(z_{2\ell}, z_{2\ell+1}) + (a_2 + a_5 K) d(z_{2\ell+1}, z_{2\ell+2}). \end{aligned} \quad (3.37)$$

Therefore,

$$d(z_{2\ell+1}, z_{2\ell+2}) \leq \frac{a_1 + a_3 + a_5 K}{1 - a_2 - a_5 K} d(z_{2\ell}, z_{2\ell+1}), \quad (3.38)$$

that is, inequality (3.34) holds in this case.

Similarly, for  $n = 2\ell$ ,  $\ell \in \mathbb{N}_0$ , we have  $d(z_{2\ell}, z_{2\ell+1}) = d(Sx_{2\ell}, Tx_{2\ell+1})$ , and from (3.33) we have

$$\begin{aligned} d(Sx_{2\ell}, Tx_{2\ell+1}) &\leq a_1 d(Fx_{2\ell}, Gx_{2\ell+1}) + a_2 d(Fx_{2\ell}, Sx_{2\ell}) \\ &\quad + a_3 d(Gx_{2\ell+1}, Tx_{2\ell+1}) + a_4 d(Fx_{2\ell}, Tx_{2\ell+1}) \\ &\quad + a_5 d(Gx_{2\ell+1}, Sx_{2\ell}), \end{aligned} \quad (3.39)$$

that is,

$$\begin{aligned} d(z_{2\ell}, z_{2\ell+1}) &\leq a_1 d(z_{2\ell-1}, z_{2\ell}) + a_2 d(z_{2\ell-1}, z_{2\ell}) + a_3 d(z_{2\ell}, z_{2\ell+1}) \\ &\quad + a_4 d(z_{2\ell-1}, z_{2\ell+1}) + a_5 d(z_{2\ell}, z_{2\ell}) \\ &= (a_1 + a_2) d(z_{2\ell-1}, z_{2\ell}) + a_3 d(z_{2\ell}, z_{2\ell+1}) + a_4 d(z_{2\ell-1}, z_{2\ell+1}) \\ &\leq (a_1 + a_2) d(z_{2\ell-1}, z_{2\ell}) + a_3 d(z_{2\ell}, z_{2\ell+1}) + a_4 K d(z_{2\ell-1}, z_{2\ell}) + a_4 K d(z_{2\ell}, z_{2\ell+1}) \\ &= (a_1 + a_2 + a_4 K) d(z_{2\ell-1}, z_{2\ell}) + (a_3 + a_4 K) d(z_{2\ell}, z_{2\ell+1}). \end{aligned} \quad (3.40)$$

Thus,

$$d(z_{2\ell}, z_{2\ell+1}) \leq \frac{a_1 + a_2 + a_4 K}{1 - a_3 - a_4 K} d(z_{2\ell-1}, z_{2\ell}), \quad (3.41)$$

and inequality (3.34) holds in this case, too.

By the same arguments as in Theorem 3.1 we conclude that  $\{z_n\}$  is a Cauchy sequence.

Let us suppose that  $SX$  is complete subspace of  $X$ . Completeness of  $SX$  implies existence of  $z \in SX$  such that  $\lim_{n \rightarrow \infty} z_{2n} = \lim_{n \rightarrow \infty} Sx_{2n} = z$ . Then, we have

$$\lim_{n \rightarrow \infty} Gx_{2n+1} = \lim_{n \rightarrow \infty} Sx_{2n} = \lim_{n \rightarrow \infty} Fx_{2n} = \lim_{n \rightarrow \infty} Tx_{2n+1} = z, \quad (3.42)$$

that is, for any  $\theta \ll c$ , for sufficiently large  $n$  we have  $d(z_n, z) \ll c$ . Since  $z \in SX \subset GX$ , there exists  $y \in X$  such that  $z = Gy$ . Let us prove that  $z = Ty$ . From  $(d_3)$  and (3.33), we have

$$\begin{aligned} d(Ty, z) &\leq Kd(Ty, Sx_{2n}) + Kd(Sx_{2n}, z) \\ &\leq a_1 Kd(Fx_{2n}, Gy) + a_2 Kd(Fx_{2n}, Sx_{2n}) + a_3 Kd(Gy, Ty) \\ &\quad + a_4 Kd(Fx_{2n}, Ty) + a_5 Kd(Gy, Sx_{2n}) + Kd(Sx_{2n}, z) \\ &= a_1 Kd(z_{2n-1}, z) + a_2 Kd(z_{2n-1}, z_{2n}) + a_3 Kd(z, Ty) \\ &\quad + a_4 Kd(z_{2n-1}, Ty) + a_5 Kd(z, z_{2n}) + Kd(z_{2n}, z) \\ &\leq a_1 Kd(z_{2n-1}, z) + a_2 Kd(z_{2n-1}, z_{2n}) + a_3 Kd(z, Ty) \\ &\quad + a_4 K^2 d(z_{2n-1}, z) + a_4 K^2 d(z, Ty) + a_5 Kd(z, z_{2n}) + Kd(z_{2n}, z). \end{aligned} \quad (3.43)$$

The sequence  $\{z_n\}$  converges to  $z$ , so for each  $c \in \text{int } P$  there exists  $n_0 \in \mathbb{N}$  such that for every  $n > n_0$

$$\begin{aligned} d(Ty, z) &\leq \frac{1}{1 - a_3 K - a_4 K^2} \\ &\quad \left( a_1 Kd(z_{2n-1}, z) + a_2 Kd(z_{2n-1}, z_{2n}) + a_4 K^2 d(z_{2n-1}, z) + a_5 Kd(z, z_{2n}) + Kd(z_{2n}, z) \right) \\ &\ll \frac{a_1 K}{1 - a_3 K - a_4 K^2} \cdot \frac{1 - a_3 K - a_4 K^2}{a_1 K} \cdot \frac{c}{5} + \frac{a_2 K}{1 - a_3 K - a_4 K^2} \cdot \frac{1 - a_3 K - a_4 K^2}{a_2 K} \cdot \frac{c}{5} \\ &\quad + \frac{a_4 K^2}{1 - a_3 K - a_4 K^2} \cdot \frac{1 - a_3 K - a_4 K^2}{a_4 K^2} \cdot \frac{c}{5} + \frac{a_5 K}{1 - a_3 K - a_4 K^2} \cdot \frac{1 - a_3 K - a_4 K^2}{a_5 K} \cdot \frac{c}{5} \\ &\quad + \frac{K}{1 - a_3 K - a_4 K^2} \cdot \frac{1 - a_3 K - a_4 K^2}{K} \cdot \frac{c}{5} \\ &= c, \end{aligned} \quad (3.44)$$

because of (3.32). Now, by  $(p_2)$  it follows that  $d(Ty, z) = \theta$ , that is,  $Ty = z$ . So, we have  $Ty = Gy = z$ , that is,  $y$  is a coincidence point, and  $z$  is a point of coincidence of mappings  $T$  and  $G$ .

Since  $TX \subset FX$ , there exists  $v \in X$  such that  $z = Fv$ . Let us prove that  $Sv = z$ , too. From  $(d_3)$  and (3.33), we have

$$\begin{aligned}
d(Sv, z) &\leq Kd(Sv, Tx_{2n+1}) + Kd(Tx_{2n+1}, z) \\
&\leq a_1Kd(Fv, Gx_{2n+1}) + a_2Kd(Fv, Sv) + a_3Kd(Gx_{2n+1}, Tx_{2n+1}) \\
&\quad + a_4Kd(Fv, Tx_{2n+1}) + a_5Kd(Gx_{2n+1}, Sv) + Kd(Tx_{2n+1}, z) \\
&= a_1Kd(z, z_{2n}) + a_2Kd(z, Sv) + a_3Kd(z_{2n}, z_{2n+1}) \\
&\quad + a_4Kd(z, z_{2n+1}) + a_5Kd(z_{2n}, Sv) + Kd(Tx_{2n+1}, z) \\
&\leq a_1Kd(z, z_{2n}) + a_2Kd(z, Sv) + a_3Kd(z_{2n}, z_{2n+1}) \\
&\quad + a_4Kd(z, z_{2n+1}) + a_5K^2d(z_{2n}, z) + a_5K^2d(Sv, z) + Kd(Tx_{2n+1}, z),
\end{aligned} \tag{3.45}$$

and by the same arguments as above, we conclude that  $d(Sv, z) = \theta$ , that is,  $Sv = Fv = z$ . Thus,  $z$  is a point of coincidence of mappings  $S$  and  $F$ , too.

Suppose that there exists  $z^*$  which is also a point of coincidence of these four mappings, that is,  $Fv^* = Gy^* = Sv^* = Ty^* = z^*$ . From (3.33) we have

$$\begin{aligned}
d(z, z^*) &= d(Sv, Ty^*) \\
&\leq a_1Kd(Fv, Gy^*) + a_2Kd(Fv, Sv) + a_3Kd(Gy^*, Ty^*) \\
&\quad + a_4Kd(Fv, Ty^*) + a_4Kd(Gy^*, Sv) \\
&= a_1Kd(z, z^*) + a_2Kd(z, z) + a_3Kd(z^*, z^*) + a_4Kd(z, z^*) + a_5Kd(z^*, z) \\
&= (a_1 + a_4 + a_5)Kd(z, z^*),
\end{aligned} \tag{3.46}$$

and (because of  $(p_3)$ ) it follows that  $z = z^*$ . Therefore,  $z$  is the unique point of coincidence of pairs  $\{S, F\}$  and  $\{T, G\}$ , and we have  $z = Sv = Fv = Gy = Ty$ . If  $\{S, F\}$  and  $\{T, G\}$  are weakly compatible pairs, then  $z$  is the unique common fixed point of  $S, F, T$ , and  $G$ , by Lemma 2.8.

The proofs for the cases in which  $FX, GX$ , or  $TX$  is complete are similar.  $\square$

Theorem 3.8 is a generalization of [13, Theorem 2.8]. Choosing  $K = 1$  from Theorem 3.8 we get the following corollary.

**Corollary 3.9.** *Let  $(X, d)$  be cone metric space and  $P$  a solid cone. Suppose that self-mappings  $F, G, S, T : X \rightarrow X$  are such that  $SX \subset GX, TX \subset FX$  and that there exist nonnegative constants  $a_i, i = 1, \dots, 5$ , satisfying  $a_1 + a_2 + a_3 + 2 \max\{a_4, a_5\} < 1$ , such that for all  $x, y \in X$  inequality*

$$d(Sx, Ty) \leq a_1d(Fx, Gy) + a_2d(Fx, Sx) + a_3d(Gy, Ty) + a_4d(Fx, Ty) + a_5d(Gy, Sx), \tag{3.47}$$

holds. If one of  $SX$ ,  $TX$ ,  $FX$ , or  $GX$  is complete subspace of  $X$ , then  $\{S, F\}$  and  $\{T, G\}$  have a unique point of coincidence in  $X$ . Moreover, if  $\{S, F\}$  and  $\{T, G\}$  are weakly compatible pairs, then  $F$ ,  $G$ ,  $S$ , and  $T$  have a unique common fixed point.

If we choose  $T = S$  and  $G = F$ , from Theorem 3.8, we get the following result for two mappings on CMTS.

**Corollary 3.10.** *Let  $(X, d)$  be a CMTS with constant  $K \geq 1$  and  $P$  a solid cone. Suppose that self-mappings  $F, S : X \rightarrow X$  are such that  $SX \subset FX$  and that there exist nonnegative constants  $a_i$ ,  $i = 1, \dots, 5$ , satisfying*

$$a_1 + a_2 + a_3 + 2K \max\{a_4, a_5\} < 1, \quad a_3K + a_4K^2 < 1, \quad a_2K + a_5K^2 < 1, \quad (3.48)$$

such that for all  $x, y \in X$  inequality

$$d(Sx, Sy) \leq a_1d(Fx, Fy) + a_2d(Fx, Sx) + a_3d(Fy, Sy) + a_4d(Fx, Sy) + a_5d(Fy, Sx), \quad (3.49)$$

holds. If one of  $SX$  or  $FX$  is complete subspace of  $X$ , then  $S$  and  $F$  have a unique point of coincidence in  $X$ . Moreover, if  $\{F, S\}$  is a weakly compatible pair, then  $F$  and  $S$  have a unique common fixed point.

## Acknowledgments

The authors are indebted to the referees for their valuable suggestions, which have contributed to improve the presentation of the paper. The first two authors were supported in part by the Serbian Ministry of Science and Technological Developments (Grant no. 174015).

## References

- [1] L.-G. Huang and X. Zhang, "Cone metric spaces and fixed point theorems of contractive mappings," *Journal of Mathematical Analysis and Applications*, vol. 332, no. 2, pp. 1468–1476, 2007.
- [2] K. Deimling, *Nonlinear Functional Analysis*, Springer, Berlin, Germany, 1985.
- [3] P. Vetro, "Common fixed points in cone metric spaces," *Rendiconti del Circolo Matematico di Palermo*, vol. 56, no. 3, pp. 464–468, 2007.
- [4] D. Ilić and V. Rakočević, "Common fixed points for maps on cone metric space," *Journal of Mathematical Analysis and Applications*, vol. 341, no. 2, pp. 876–882, 2008.
- [5] M. Abbas and G. Jungck, "Common fixed point results for noncommuting mappings without continuity in cone metric spaces," *Journal of Mathematical Analysis and Applications*, vol. 341, no. 1, pp. 416–420, 2008.
- [6] C. Di Bari and P. Vetro, " $\varphi$ -pairs and common fixed points in cone metric spaces," *Rendiconti del Circolo Matematico di Palermo*, vol. 57, no. 2, pp. 279–285, 2008.
- [7] S. Radenović, "Common fixed points under contractive conditions in cone metric spaces," *Computers & Mathematics with Applications*, vol. 58, no. 6, pp. 1273–1278, 2009.
- [8] G. Jungck, S. Radenović, S. Radojević, and V. Rakočević, "Common fixed point theorems for weakly compatible pairs on cone metric spaces," *Fixed Point Theory and Applications*, vol. 2009, Article ID 643840, 13 pages, 2009.
- [9] B. S. Choudhury and N. Metiya, "The point of coincidence and common fixed point for a pair of mappings in cone metric spaces," *Computers & Mathematics with Applications*, vol. 60, no. 6, pp. 1686–1695, 2010.

- [10] F. Sabetghadam and H. P. Masiha, "Common fixed points for generalized  $\varphi$ -pair mappings on cone metric spaces," *Fixed Point Theory and Applications*, vol. 2010, Article ID 718340, 8 pages, 2010.
- [11] B. Fisher, "Four mappings with a common fixed point," *The Journal of the University of Kuwait. Science*, vol. 8, pp. 131–139, 1981.
- [12] Z. Kadelburg, S. Radenović, and V. Rakočević, "Topological vector space-valued cone metric spaces and fixed point theorems," *Fixed Point Theory and Applications*, vol. 2010, Article ID 170253, 17 pages, 2010.
- [13] M. Abbas, B. E. Rhoades, and T. Nazir, "Common fixed points for four maps in cone metric spaces," *Applied Mathematics and Computation*, vol. 216, no. 1, pp. 80–86, 2010.
- [14] S. Janković, Z. Golubović, and S. Radenović, "Compatible and weakly compatible mappings in cone metric spaces," *Mathematical and Computer Modelling*, vol. 52, no. 9-10, pp. 1728–1738, 2010.
- [15] Sh. Rezapour and R. Hamlbarani, "Some notes on the paper: "Cone metric spaces and fixed point theorems of contractive mappings"," *Journal of Mathematical Analysis and Applications*, vol. 345, no. 2, pp. 719–724, 2008.
- [16] M. A. Khamsi and N. Hussain, "KKM mappings in metric type spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 73, no. 9, pp. 3123–3129, 2010.
- [17] G. Jungck and B. E. Rhoades, "Fixed point theorems for occasionally weakly compatible mappings," *Fixed Point Theory*, vol. 7, no. 2, pp. 287–296, 2006.
- [18] D. S. Mitrinović, *Analytic Inequalities*, vol. 165 of *Grundlehren der mathematischen Wissenschaften*, Springer, New York, NY, USA, 1970.